Theoretical 3: Solution

Physics of Spin

Part A. Larmor Precession

1. From the two equations given in the text, we obtain the relation

$$\frac{d\boldsymbol{\mu}}{dt} = -\gamma\boldsymbol{\mu} \times \mathbf{B}. \tag{1}$$

Taking the dot product of eq (1). with $\boldsymbol{\mu}$, we can prove that

$$\begin{aligned} \boldsymbol{\mu} \cdot \frac{d\boldsymbol{\mu}}{dt} &= -\gamma\boldsymbol{\mu} \cdot (\boldsymbol{\mu} \times \mathbf{B}), \\ \frac{d\left|\boldsymbol{\mu}\right|^2}{dt} &= 0, \\ \mu = \left|\boldsymbol{\mu}\right| &= \text{const.} \end{aligned} \tag{2}$$

Taking the dot product of eq. (1) with $\mathbf{B}$, we also prove that

$$\begin{aligned} \mathbf{B} \cdot \frac{d\boldsymbol{\mu}}{dt} &= -\gamma\mathbf{B} \cdot (\boldsymbol{\mu} \times \mathbf{B}), \\ \mathbf{B} \cdot \frac{d\boldsymbol{\mu}}{dt} &= 0, \\ \mathbf{B} \cdot \boldsymbol{\mu} &= \text{const.} \end{aligned} \tag{3}$$

An acute reader will notice that our master equation in (1) is identical to the equation of motion for a charged particle in a magnetic field

$$\frac{d\mathbf{v}}{dt} = \frac{q}{m}\mathbf{v} \times \mathbf{B}. \tag{4}$$

Hence, the same argument for a charged particle in magnetic field can be applied in this case.

2. For a magnetic moment making an angle of ϕ with $\mathbf{B}$,

$$\begin{aligned} \frac{d\boldsymbol{\mu}}{dt} &= -\gamma\boldsymbol{\mu} \times \mathbf{B}, \\ \left|\boldsymbol{\mu}\right| \sin\phi \frac{d\theta}{dt} &= \gamma \left|\boldsymbol{\mu}\right| B_0 \sin\phi, \\ \omega_0 = \frac{d\theta}{dt} &= \gamma B_0. \end{aligned} \tag{5}$$

Part B. Rotating frame

1. Using the relation given in the text, it is easily shown that

$$\begin{aligned} \left(\frac{d\boldsymbol{\mu}}{dt}\right)_{\text{rot}} &= \left(\frac{d\boldsymbol{\mu}}{dt}\right)_{\text{lab}} - \boldsymbol{\omega} \times \boldsymbol{\mu} \\ &= -\gamma\boldsymbol{\mu} \times \mathbf{B} - \omega\mathbf{k}' \times \boldsymbol{\mu} \\ &= -\gamma\boldsymbol{\mu} \times \left(\mathbf{B} - \frac{\omega}{\gamma}\mathbf{k}'\right) \\ &= -\gamma\boldsymbol{\mu} \times \mathbf{B}_{\text{eff}}. \end{aligned} \tag{6}$$

Note that $\mathbf{k}$ is equal to $\mathbf{k}'$ as observed in the rotating frame.

2. The new precession frequency as viewed on the rotating frame S' is

$$\begin{aligned} \vec{\Delta} &= (\omega_0 - \omega)\,\mathbf{k}', \\ \Delta &= \gamma B_0 - \omega. \end{aligned} \tag{7}$$

3. Since the magnetic field as viewed in the rotating frame is $\mathbf{B} = B_0\mathbf{k}' + b\mathbf{i}'$,

$$\mathbf{B}_{\text{eff}} = \mathbf{B} - \omega/\gamma\mathbf{k}' = \left(B_0 - \frac{\omega}{\gamma}\right)\mathbf{k}' + b\mathbf{i}',$$

and

$$\begin{aligned}\Omega &= \gamma \left|\mathbf{B}_{\text{eff}}\right|, \\ &= \gamma\sqrt{\left(B_0 - \frac{\omega}{\gamma}\right)^2 + b^2}.\end{aligned} \tag{8}$$

4. In this case, the effective magnetic field becomes

$$\begin{aligned}\mathbf{B}_{\text{eff}} &= \mathbf{B} - \omega/\gamma\mathbf{k}' \\ &= \left(B_0 - \frac{\omega}{\gamma}\right)\mathbf{k}' + b(\cos 2\omega t\mathbf{i}' - \sin 2\omega t\mathbf{j}')\end{aligned} \tag{9}$$

which has a time average of $\overline{\mathbf{B}_{\text{eff}}} = \left(B_0 - \frac{\omega}{\gamma}\right)\mathbf{k}'$.

Part C. Rabi oscillation

1. The oscillating field can be considered as a superposition of two oppositely rotating field:

$$2b\cos\omega_0 t\mathbf{i} = b\left(\cos\omega_0 t\mathbf{i} + \sin\omega_0 t\mathbf{j}\right) + b\left(\cos\omega_0 t\mathbf{i} - \sin\omega_0 t\mathbf{j}\right),$$

which gives an effective field of (with $\omega = \omega_0 = \gamma B_0$):

$$\mathbf{B}_{\text{eff}} = \left(B_0 - \frac{\omega}{\gamma}\right)\mathbf{k}' + b\mathbf{i}' + b(\cos 2\omega_0 t\mathbf{i}' - \sin 2\omega_0 t\mathbf{j}').$$

Since $\omega_0 \gg \gamma b$, the rotation of the term $b(\cos 2\omega_0 t\mathbf{i}' - \sin 2\omega_0 t\mathbf{j}')$ is so fast compared to the frequency γb. This means that we can take the approximation


$$\mathbf{B}_{\text{eff}} \approx \left(B_0 - \frac{\omega}{\gamma}\right)\mathbf{k}' + b\mathbf{i}' = b\mathbf{i}', \tag{10}$$

where the magnetic moment precesses with frequency $\Omega = \gamma b$.

As $\Omega = \gamma b \ll \omega_0$, the magnetic moment does not "feel" the rotating term $b\left(\cos 2\omega_0 t\mathbf{i}' - \sin 2\omega\right.$ which averaged to zero.

2. Since the angle α that $\boldsymbol{\mu}$ makes with $\mathbf{B}_{\text{eff}}$ stays constant and $\boldsymbol{\mu}$ is initially oriented along the z axis, α is also the angle between $\mathbf{B}_{\text{eff}}$ and the z axis which is

$$\tan\alpha = \frac{b}{B_0 - \frac{\omega}{\gamma}}. \tag{11}$$



From the geometry of the system, we can show that ($\cos\theta = \mu_z/\mu$):

$$\begin{aligned}2\mu\sin\frac{\theta}{2} &= 2\mu\sin\alpha\sin\frac{\Omega t}{2}, \\ \sin^2\frac{\theta}{2} &= \sin^2\alpha\sin^2\frac{\Omega t}{2}, \\ \frac{1-\cos\theta}{2} &= \sin^2\alpha\frac{1-\cos\Omega t}{2}, \\ \cos\theta &= 1 - \sin^2\alpha + \sin^2\alpha\cos\Omega t, \\ \cos\theta &= \cos^2\alpha + \sin^2\alpha\cos\Omega t.\end{aligned}$$

So, the projected magnetic moment along the z axis is $\mu_z(t) = \mu\cos\theta$ and the magnetization is

$$M = N\mu_z = N\mu\left(\cos^2\alpha + \sin^2\alpha\cos\Omega t\right). \tag{12}$$

Note that the magnetization does not depend on the reference frame S or S' (μ_z has the same value viewed in both frames).

Taking $\omega = \omega_0 = \gamma B_0$, the angle α is 90^0 and $M = N\mu\cos\Omega t$.

3. From the relations


$$P_\uparrow - P_\downarrow = \frac{\mu_z}{\mu} = \cos\theta,$$
$$P_\uparrow + P_\downarrow = 1,$$

we obtain the results ($\omega = \omega_0$)

$$\begin{aligned} P_\downarrow &= \frac{1-\cos\theta}{2} \\ &= \frac{1-\cos^2\alpha - \sin^2\alpha\cos\Omega t}{2} \\ &= \sin^2\alpha\frac{1-\cos\Omega t}{2} \\ &= \frac{b^2}{\left(B_0 - \frac{\omega}{\gamma}\right)^2 + b^2}\sin^2\frac{\Omega t}{2} \\ &= \sin^2\frac{\Omega t}{2}, \end{aligned} \tag{13}$$

and

$$P_\uparrow = \frac{b^2}{\left(B_0 - \frac{\omega}{\gamma}\right)^2 + b^2}\cos^2\frac{\Omega t}{2} = \cos^2\frac{\Omega t}{2}. \tag{14}$$



Part D. Measurement incompatibility

1. In the x direction, the uncertainty in position due to the screen opening is Δx. According to the uncertainty principle, the atom momentum uncertainty Δp_x is given by

$$\Delta p_x \approx \frac{\hbar}{\Delta x},$$

which translates into an uncertainty in the x velocity of the atom,

$$v_x \approx \frac{\hbar}{m\Delta x}.$$

Consequently, during the time of flight t of the atoms through the device, the uncertainty in the width of the beam will grow by an amount δx given by

$$\delta x = \Delta v_x t \approx \frac{\hbar}{m\Delta x}t.$$

So, the width of the beams is growing linearly in time. Meanwhile, the two beams are separating at a rate determined by the force F_x and the separation between the beams after a time t becomes

$$d_x = 2\times\frac{1}{2}\frac{F_x}{m}t^2 = \frac{1}{m}\left|\mu_x\right|Ct^2.$$

In order to be able to distinguish which beam a particle belongs to, the separation of the two beams must be greater than the widths of the beams; otherwise the two beams will overlap and it will be impossible to know what the x component of the atom spin is. Thus, the condition must be satisfied is

$$\begin{aligned} d_x &\gg \delta x, \\ \frac{1}{m}\left|\mu_x\right| Ct^2 &\gg \frac{\hbar}{m\Delta x}t, \\ \frac{1}{\hbar}\left|\mu_x\right| \Delta x Ct &\gg 1. \end{aligned} \tag{15}$$

2. As the atoms pass through the screen, the variation of magnetic field strength across the beam width experienced by the atoms is

$$\Delta B = \Delta x \frac{dB}{dx} = C\Delta x.$$

This means the atoms will precess at rates covering a range of values $\Delta\omega$ given by

$$\Delta\omega = \gamma\Delta B = \frac{\mu_z}{\hbar}\Delta B = \frac{|\mu_x|}{\hbar}C\Delta x,$$

and, if previous condition in measuring μ_x is satisfied,

$$\Delta\omega t \gg 1. \tag{16}$$

In other words, the spread in the angle $\Delta\omega t$ through which the magnetic moments precess is so large that the z component of the spin is completely randomized or the measurement uncertainty is very large.