Particles and Waves (10 points)

Part A. Quantum particle in a box (1.4 points)

A.1 (0.4 points)

The width of the potential well (L) should be equal to the half of the wavelength of the de Broglie standing wave $\lambda_{\mathrm{dB}} = h/p$, here h is the Planck's constant and p is the momentum of the particle. Thus $p = h/\lambda_{\mathrm{dB}} = h/(2L)$, and the minimal possible energy of the particle is

$$E_{\min} = \frac{p^2}{2m} = \frac{h^2}{8mL^2}.$$

A.2 (0.6 points)

The potential well should fit an integer number of the de Broglie half-wavelengths: $L = \frac{1}{2}\lambda_{\mathrm{dB}}^{(n)} \cdot n$, $n = 1, 2, \ldots$. Therefore, particle's momentum, corresponding to the de Broglie wavelength $\lambda_{\mathrm{dB}}^{(n)}$ is

$$p_n = \frac{h}{\lambda_{\mathrm{dB}}^{(n)}} = \frac{hn}{2L},$$

and the corresponding energy is

$$E_n = \frac{p_n^2}{2m} = \frac{h^2 n^2}{8mL^2}, \qquad n = 1, 2, 3, \ldots. \tag{1}$$

A.3 (0.4 points)

The energy of the emitted photon, $E = hc/\lambda$ (here c is the speed of light and λ is the photon's wavelength) should be equal to the energy difference $\Delta E = E_2 - E_1$, therefore

$$\lambda_{21} = \frac{hc}{E_2 - E_1} = \frac{8mcL^2}{3h}.$$

Part B. Optical properties of molecules (2.1 points)

B.1 (0.8 points)

Taking into account the Pauli exclusion principle, each energy level E_n is occupied by two electrons with spins oriented in the opposite directions. As a results, 10 electrons fill the lowest 5 states, and the absorption of the photon of the longest wavelength corresponds to the transition of one electron from the occupied E_5 to the unoccupied E_6 energy state:

$$\frac{hc}{\lambda} = E_6 - E_5,$$

where E_6 and E_5 can be found from Eq. 1, where m is replaced with the electron mass m_{e}. Hence we obtain:

$$\lambda = \frac{c \cdot 8m_{\mathrm{e}}L^2}{h\,(6^2 - 5^2)} = \frac{10.5^2 \cdot 8}{11}\,\frac{m_{\mathrm{e}}cl^2}{h} = \frac{882}{11}\,\frac{m_{\mathrm{e}}cl^2}{h} \approx 647\,\mathrm{nm}.$$

This result correspond precisely to the experimental value the peak position of the Cy5 absorption spectrum.

B.2 (0.4 points)

In the similar model for the Cy3 molecule, there are 8 electrons in the box of length $L = 8.5l$, thus photon's absorption corresponds to the $E_4 \to E_5$ transition. Taking into account the result of question B1, we obtain

$$\lambda_{\mathrm{Cy3}} = \frac{8.5^2 \cdot 8}{(5^2 - 4^2)}\,\frac{m_{\mathrm{e}}cl^2}{h} \approx 518\,\mathrm{nm},$$

i. e. the absorption spectrum of the Cy3 molecule is shifted by $\Delta\lambda \approx 129\,\mathrm{nm}$ to the blue comparing to that of the Cy5 molecule. The experimental value is $\lambda_{\mathrm{Cy3}}^{(\mathrm{exp})} = 548\,\mathrm{nm}$, so that our model catches general properties of these dye molecules rather well.

B.3 (0.7 points)

Let us assume

$$K = k\varepsilon_0^{\alpha} h^{\beta} \lambda^{\gamma} d^{\delta}. \tag{2}$$

The SI units of the relevant quantities are:

$$[\varepsilon_0] = \frac{\mathrm{A}^2 \cdot \mathrm{s}^4}{\mathrm{kg} \cdot \mathrm{m}^3}, \qquad [h] = \frac{\mathrm{kg} \cdot \mathrm{m}^2}{\mathrm{s}}, \qquad [\lambda] = \mathrm{m}, \qquad [d] = \mathrm{A} \cdot \mathrm{s} \cdot \mathrm{m}, \qquad [K] = \mathrm{s}^{-1}.$$

By plugging these expressions into Eq. 2 we obtain a simple system of linear equations for the unknown powers α, β, γ, and δ:

$$2\alpha + \delta = 0, \qquad -\alpha + \beta = 0, \qquad 4\alpha - \beta + \delta = -1, \qquad -3\alpha + 2\beta + \gamma + \delta = 0.$$

By solving this system we get:

$$\alpha = \beta = -1, \qquad \gamma = -3, \qquad \delta = 2,$$

so that the rate of spontaneous emission is

$$K = \frac{16\pi^3}{3} \frac{d^2}{\varepsilon_0 h \lambda^3}. \tag{3}$$

B.4 (0.2 points)

By using the result of question B.2 and expressing transition dipole moment as $d = 2.4\, el$, we obtain from Eq. 3:

$$\tau_{\mathrm{Cy5}} = \frac{3}{16\pi^3} \frac{\varepsilon_0 h}{2.4^2 l^2 e^2} \lambda^3 \approx 3.3\,\mathrm{ns}.$$

Part C. Bose-Einstein condensation (1.5 points)

C.1 (0.4 points)

At temperature T, the average kinetic energy of translational motion is $\frac{3}{2} k_{\mathrm{B}} T$. Equating this result to $p^2/(2m)$, we obtain typical momentum $p = \sqrt{3mk_{\mathrm{B}}T}$ and the de Broglie wavelength

$$\lambda_{\mathrm{dB}} = \frac{h}{p} = \frac{h}{\sqrt{3mk_{\mathrm{B}}T}}.$$

C.2 (0.5 points)

The volume per particle V/N is a good estimate for ℓ^3. We obtain $\ell = n^{-1/3}$, with $n = N/V$ and equate $\ell = \lambda_{\mathrm{dB}}$ to express $T_c = h^2 n^{2/3}/(3mk_{\mathrm{B}})$.

C.3 (0.6 points)

Using the answer to the previous question, we express $n_c = (3mk_{\mathrm{B}}T_c)^{3/2}/h^3$. Equation of state for the ideal gas gives $n_0 = p/(k_{\mathrm{B}}T)$. Numerical estimations yield $n_c \approx 1.59 \cdot 10^{18}\,\mathrm{m}^{-3}$ and $n_0/n_c \approx 1.5 \cdot 10^7$.

Part D. Three-beam optical lattices (5.0 points)

D.1 (1.4 points)

We sum the three electric fields (z components)

$$E(\vec{r}, t) = E_0 \sum_{i=1}^{3} \cos\left(\vec{k}_i \cdot \vec{r} - \omega t\right), \tag{4}$$

and square the result

$$\begin{aligned} E^2(\vec{r}, t) &= E_0^2 \sum_{i=1}^{3} \sum_{j=1}^{3} \cos\left(\vec{k}_i \cdot \vec{r} - \omega t\right) \cos\left(\vec{k}_j \cdot \vec{r} - \omega t\right) \\ &= \frac{E_0^2}{2} \sum_{i=1}^{3} \sum_{j=1}^{3} \left\{ \cos\left[\left(\vec{k}_i - \vec{k}_j\right) \cdot \vec{r}\right] + \cos\left[\left(\vec{k}_i + \vec{k}_j\right) \cdot \vec{r} - 2\omega t\right] \right\}. \end{aligned} \tag{5}$$

Time averaging gives

$$\langle E^2(\vec{r},t)\rangle = \frac{E_0^2}{2}\sum_{i=1}^{3}\sum_{j=1}^{3}\cos\left[\left(\vec{k}_i-\vec{k}_j\right)\cdot\vec{r}\right], \tag{6}$$

we analyse the 9 terms and simplify to

$$\langle E^2(\vec{r},t)\rangle = E_0^2\left(\frac{3}{2}+\sum_{j=1}^{3}\cos\left(\vec{b}_j\cdot\vec{r}\right)\right). \tag{7}$$

Here

$$\vec{b}_1=\vec{k}_2-\vec{k}_3, \qquad \vec{b}_2=\vec{k}_3-\vec{k}_1, \qquad \vec{b}_3=\vec{k}_1-\vec{k}_2,$$

or in terms of the Levi-Civita symbol, $\vec{b}_k=\varepsilon_{ijk}(\vec{k}_i-\vec{k}_j)$. Incidentally, they are known as the reciprocal lattice vectors.

D.2 (0.5 points)

Argument: Observe that rotation by 60° maps the three vectors $\vec{b}_{1,2,3}$ into the relabelled triplet of $-\vec{b}$'s.

D.3 (1.2 points)

We find

$$V(x,y)=-\alpha E_0^2\left\{\frac{3}{2}+\cos\left(ky\sqrt{3}\right)+\cos\left(\frac{3kx}{2}+\frac{ky\sqrt{3}}{2}\right)+\cos\left(\frac{3kx}{2}-\frac{ky\sqrt{3}}{2}\right)\right\}, \tag{8}$$

and deduce

$$V_X(x)=-\alpha E_0^2\left\{\frac{5}{2}+2\cos\frac{3kx}{2}\right\}. \tag{9}$$

The potential has a simple cosine form, and the origin in an obvious minimum. Its replica appear at multiples of $\Delta x=4\pi/(3k)$. In the midpoint between any two minima, e.g. at $x=\Delta x/2=2\pi/(3k)$, the function $V_X(x)$ has its maxima.

Concerning the behaviour along the y axis, we have

$$V_Y(y)=-\alpha E_0^2\left\{\frac{3}{2}+\cos 2\varphi+2\cos\varphi\right\}, \qquad \varphi=\sqrt{3}ky/2. \tag{10}$$

Looking for the extrema, we find the equation

$$\sin 2\varphi+\sin\varphi=0. \tag{11}$$

∘ $\varphi=0$ (corresponding to $y=0$) is the 'deep' minimum – the lattice site;

∘ $\varphi=\pi$ (corresponding to $y=\frac{2\pi}{\sqrt{3}k}$) is the 'shallow' minimum (later shown to be a saddle point of $V(x,y)$);

∘ $\varphi=2\pi/3$ and $\varphi=4\pi/3$ (corresponding to $y=\frac{4\pi}{3\sqrt{3}k}$ and $y=\frac{8\pi}{3\sqrt{3}k}$, respectively) are maxima.

D.4 (0.8 points)

We review the minima found in the previous question and eliminate the saddle point at $(0,2\pi/3\sqrt{3}\,k)$. The actual minima of the 2D potential landscape $V(x,y)$ are:

∘ $(0,0)$ – at the origin;

∘ $(4\pi/(3k),0)$ – nearest to the origin in the positive direction along the x axis. On the grounds of symmetry we argue that there are six equivalent nearest minima in the directions 0°, ±60°, ±120°, and 180° with respect to the x axis.

Distance between nearest minima (the lattice constant) $a=4\pi/(3k)$. Given that the laser wavelength is $\lambda_{\rm las}=2\pi/k$, we have $a=\Delta x=2\lambda_{\rm las}/3$, thus $a/\lambda_{\rm las}=2/3$.

D.5 (1.1 points)

The atom's core electrons (all but the one promoted to to a state with a high principal quantum number n) shield the electric field of the nucleus so that the effective potential for the outer electron resembles that of a hydrogen atom. The attractive force acting on that electron, $F = e^2/(4\pi\varepsilon_0 r^2)$, gives rise to its centripetal acceleration $a = v^2/r$. Equating $F = m_\mathrm{e}a$ and using the expression for the angular momentum $m_\mathrm{e}vr = n\hbar$ to eliminate the velocity, we find the quantum number n corresponding to the orbit with the radius $r = \lambda_\mathrm{las}$:

$$n = \frac{e}{\hbar}\sqrt{\frac{m_\mathrm{e}\lambda}{4\pi\varepsilon_0}} \approx 85. \tag{12}$$