

7. CHARGED ROD (6 points) — Solution by Jaan Kalda, grading schemes by...

i) (2 points) Notice that all particles with the same charge-to-mass ratio orbit in a homogeneous magnetic field B with the same frequency $\omega_B = \frac{Bq}{m}$; the orbit is a circle of radius $r = \frac{v}{\omega_B}$. Indeed, the Lorentz force must provide the centripetal acceleration, hence $Bqv = mv\omega_B \Rightarrow \omega_B = \frac{Bq}{m}$.

Since the mass-to-charge ratio is the same for all the fictitious pieces of the rod, if the rod rotates with angular speed $\omega = \frac{Bq}{m}$, then the Lorentz force provides exactly the needed acceleration to all these pieces, and the tension force is zero everywhere.

On the other hand, if the angular speed were smaller or larger, we would have either $Bqv < mv\omega$ or $Bqv > mv\omega$ for all the pieces, resulting in either stretching or compressive tension force at the centre of the rod, respectively. Therefore, the answer is $\omega = \frac{Bq}{m}$.

Grading: (preliminary)

- Considers forces on an infinitesimal part of the rod **0.4 pts**
- Equates, with justification, Lorentz and centrifugal forces $dqvB = dm\omega^2 r$ **0.4 pts**
- Uses $\omega = \frac{v}{r}$ **0.4 pts**
- Uses $\frac{dq}{dm} = \frac{q}{m}$ **0.4 pts**
- Expresses $\omega = \frac{qB}{m}$ **0.4 pts**

ii) (4 points) To begin with, let us notice that if a system of charges with the same charge-to-mass ratio $\alpha = q/m$ moves in a homogeneous magnetic field, the centre of mass will move along a circle with cyclotron frequency ω_B . Indeed, the total Lorentz force acting on the system is

$$\begin{aligned}\sum_i q_i \frac{d\vec{r}_i}{dt} \times \vec{B} &= \alpha \sum_i m_i \frac{d\vec{r}_i}{dt} \times \vec{B} \\ &= \alpha \frac{d\vec{r}_C}{dt} \times \vec{B} \sum_i m_i.\end{aligned}$$

So, Newton's second law reads

$$\frac{d^2 \vec{r}_C}{dt^2} \sum_i m_i = \frac{q}{m} \frac{d\vec{r}_C}{dt} \times \vec{B} \sum_i m_i.$$

The total mass of the system cancels out from this equation, and we can see that the centre of mass $\vec{r}_C$ moves in the same way as a point charge q with mass m .

Alternatively, the same can be achieved through integration. From Newton's second law $m\vec{a}_C = \int dq \vec{v} \times \vec{B}$, where $\vec{v}$ is the velocity of the charge element dq . But since the mass and charge distributions are homogenous, $dq = \frac{q}{m} dm$ and $\vec{B}$ is constant

so it can be taken out from the integral to achieve $m\vec{a}_C = \frac{q}{m} \left(\int \vec{v} dm \right) \times \vec{B}$. But now the integral is just $m\vec{v}_C$ (can be seen directly or through the definition of centre of mass $\int \frac{d\vec{r}}{dt} dm = \frac{d}{dt} \int \vec{r} dm = m\vec{v}_C$). Thus we get $m\vec{a}_C = q\vec{v}_C \times \vec{B}$.

Additionally, the rod can (and will) rotate with a constant speed. The fact that the angular speed must be constant follows from the conservation of kinetic energy of the rod, which is the sum of the kinetic energy of its centre of mass and the rotational energy around the centre of mass. The former is constant, so the latter must be as well.

The centre of mass moves with speed $v/2$ and draws a circle of radius $R = \frac{mv}{2Bq}$ that passes through the point $(\frac{l}{2}, 0)$ and for which the x -axis is a symmetry axis. The red end can reach the origin only when the centre of mass is at a distance $\frac{l}{2}$ from the origin. This can happen either after a full cyclotron period $T = 2\pi/\omega_B$, or at any moment assuming $R = \frac{l}{2}$ and the circle is centred around the origin.

To determine if this can happen earlier than after time T , let us assume that $R = \frac{mv}{2Bq} = \frac{l}{2}$. In that case, the angular speed of the rod's rotation is $\Omega = \frac{v}{l}$, and we get $v = \frac{Bql}{m}$. This means $\Omega = \frac{Bq}{m} = \omega_B$, i. e., the rod's rotational angular speed is the same as the centre of mass' angular speed in its orbit, which would cause the blue end to remain at the origin.

Next, we examine if the red end can reach the origin after time T . For this to happen, the condition is that $\Omega T = (2\pi n + \pi)$, where n is an integer. Since $\Omega = v/l$ and we need to minimize v , we take $n = 0$ to obtain $v = \pi l/T = \frac{lBq}{2m}$.

Grading: (preliminary)

- Deduces, with justification, that the net force on the rod $\vec{F} = q\vec{v}_C \times \vec{B}$ **0.5 pts**
- Uses $v_C = v/2$ **0.2 pts**
- Justifies that the COM moves on a circular path **0.3 pts**
- Expresses the radius of the path traced by the COM $R = \frac{mv}{2qB}$ **0.3 pts**
- Concludes that the angular velocity of the COM is $\omega = \frac{qB}{m}$ **0.2 pts**
- Expresses the angular velocity of the rotation around the COM $\Omega = \frac{v}{l}$ **0.2 pts**
- Justifies that Ω is conserved **0.3 pts**
- Argues that $t < \frac{2\pi}{\omega}$ is possible only if $R = l/2$ **0.5 pts**
- Justifies that in this case, the red end will never end up at the origin **0.5 pts**
- Justifies that $\Omega t = \pi + 2\pi k$ with $k \in \mathbb{Z}_{\geq 0}$ **0.4 pts**
- Expresses $v = \frac{qBl}{m}(\frac{1}{2} + k)$ **0.6 pts**